

CD74 Ab-1

Catalog # MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200ug/ml)

Catalog # MS-131-R7 (7.0ml)

Catalog # MS-131-PCS

INTENDED USE:

- **For In Vitro Diagnostic Use:** This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.
- **Description:** Expression of the invariant chain of the HLA-DR antigen (CD74) by B-lymphocytes begins in the pre-B stage but lost before the final differentiation to plasma cells.
- **Expected Staining Pattern:** Cell membrane
- **Positive Control:** Raji cells. Tonsil / lymph node

MATERIALS PROVIDED:

CD74 Ab-1 (refer to catalog number):

- #MS-131-P (or -P0, -P1): 200ug/ml of antibody purified from ascites. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide.
or
- #MS-131-R7: (7.0ml) of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.
or
- #MS-131-PCS: 5 positive control slides.
- **Antibody Concentration:** 200ug/ml
- **Host:** Mouse
- **Mol. Wt. of Antigen:** 35kDa
- **Epitope:** Not determined
- **Species Reactivity:** Human and Baboon. Does not react with Rat. Others-not known.
- **Clone Designation:** LN2
- **Ig Isotype / Light Chain:** IgG1 / kappa
- **Immunogen:** SU-DHL-4 lymphoma cells.
- **Microbiological State:** This product is not sterile.

MATERIALS REQUIRED, BUT NOT PROVIDED:

- **Antibody Diluent:** For concentrated antibodies, the antibody must be diluted before using. Use Lab Vision Antibody Diluent (catalog # TA-125-UD). Refer to diluent product instructions for use.
- **Negative Control Reagent:** Refer to the "General Protocol" instructions.
- **Visualization System:** Refer to the "General Protocol" instructions.

METHODS AND PROCEDURES:

Specimen Preparation	Refer to the "General Protocol" instructions.
Dilution of Concentrated Antibody	1:50-1:100 in antibody diluent
Tissue Section Pretreatment	Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0 (Lab Vision catalog # AP-9003), for 10-20 minutes followed by cooling at room temperature for 20 min.
Primary Antibody Incubation Time	30 minutes at Room Temperature
Visualization	To detect antibody, follow the instructions provided with the visualization system.

STORAGE and STABILITY:

This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

REFERENCES:

- 1) Ioachim HL; et al. International Journal of Cancer. Supplement, 1994, 8:132-3.
- 2) Kumar S; et al. American Journal of Hematology, 1994, 46(2):134-7.