

CD51 Ab-1 (Clone CJ00)

Mouse Monoclonal Antibody

Cat. #MS-1774-S0, -S1, or -S (0.1ml, 0.5ml, or 1.0ml Supernatant)

Cat. #MS-1774-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #MS-1774-PCS (5 Slides) (Positive Control for Histology)

Description: CD51 / Integrin αv is expressed as a heterodimer noncovalently associated with integrin beta3 (CD61); which is known as vitronectin receptor. CD51 is post-translationally cleaved into a large fragment of 125kDa and a small fragment of 24kDa which are disulphide linked. CD51-CD61 binds to the RGD sequence in the extracellular matrix proteins like vitronectin, von Willebrand factor, fibrinogen and thrombospondin. CD51-CD61 is expressed on platelets, endothelial cells, certain activated leukocytes, NK cells, macrophages and neutrophils.

Mol. Wt. of Antigen: 125/24kDa (reduced) / 150kDa (nonreduced)¹

Epitope: C-terminal

Ig Isotype / Light Chain: IgG₁ / k

Species Reactivity: Human. Others-not known.

Clone Designation: CJ00

Immunogen: Recombinant protein corresponding to C-terminus of human CD51

Applications and Suggested Dilutions:

- Immunohistology (Formalin/paraffin)
(Use Ab at 1:40 for 30 min at RT)
- * (Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 1mM EDTA, pH 8.0, (**NEOMARKERS'** Cat. #AP-9004), for 10-20 min followed by cooling at RT for 20 min.)

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Tonsil

Cellular Localization: Cell membrane

Supplied As: Tissue culture supernatant with 0.09% sodium azide, or Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Storage and Stability: Store vial at 4°C. When stored at 2-8°C, this antibody is stable for 24 months.

Suggested References:

1. The Leucocyte Antigen. Barclay A N, et al. Editors; Acaemic Press, Pg 218.
2. Horton M, et al. (1990) Int J Exp Pathol, 71:741-759.

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only

