

Eff. Date: 28 October 2020

Version: 1.0

IFU: Factor XIIIa ILM1448

Factor XIIIa clone F13A1/1447

Mouse Monoclonal Antibody

Instructions For Use

Specification:

It recognizes a protein of 83kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macrophages and dermal dendritic cells. Anti-factor XIIIa has been found to be useful in differentiating between dermatofibroma (almost all cases are positive), dermatofibrosarcoma protuberans (-/+) and desmoplastic malignant melanoma (-).

Catalog No.	Contents	Volume
ILM1448-C01	Factor XIIIa clone F13A1/1447	0,1 ml concentrate
ILM1448-C05	Factor XIIIa clone F13A1/1447	0,5 ml concentrate
ILM1448-C1	Factor XIIIa clone F13A1/1447	1,0 ml concentrate

Intended use: For Research Use Only **Reactivity:** Human, others not known

Clone: F13A1/1447

Species of origin: Mouse

Isotype: $IgG2a\lambda$

Human Entrez Gene ID: 2162 Human Unigene: 335513

Human Chromosome location: 6p25.3-p24.3

Control tissue: Placenta and 293T cells **Staining:** Cytoplasmic and secreted

0.05% Azide.

Immunogen: Recombinant fragment of human Factor XIIIa protein

Application and suggested dilutions:

Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, for 20 minutes is required for IHC staining on formalin-fixed, paraffin embedded tissue sections.

- Paraffin embedded tissue section, dilution up to
- ELISA
- Flow Cytometry (1-2ug/million cells)
- Immunofluorescence (11-2ug/ml)

The optimal dilution for a specific application should be determined by the investigator.

Note: Dilution of the antibody in 10% normal goat serum followed by a goat anti-Mouse secondary antibody-based detection is recommended.

Storage & Stability: Store at 2-8 °C. Do not use after expiration date printed on the vial.

References

 Davie EW. Et al. 1991. The coagulation cascade: Initiation maintenance and regulation. Biochemistry 30: 10363-10370

