

BrightVision, one component detection system Goat Anti- Mouse/Rabbit IgG HRP (Ready-to-Use)

Instruction For Use

These instructions apply to the WellMed BrightVision; one step detection system Goat Anti-Mouse/Rabbit IgG HRP (Ready-to-Use).

- 1. Intended Use
- 2. Summary and explanation
- 3. Kit components
- 4. Availability
- 5. Recommended Staining Protocol
- 6. Control slides
- 7. Storage
- 8. Warnings and precautions
- 9. Troubleshooting
- 10. Reference

1: Intended Use

For in-vitro Diagnostic Use.

WellMed BrightVision one step detection system peroxidase Goat Anti- Mouse/Rabbit IgG HRP, is intended for use in immunohistochemistry for the detection of mouse and rabbit antibodies.

2: Summary and explanation

WellMed BrightVision detection system peroxidase Goat anti- Mouse/Rabbit HRP, is a ready to use system that has been manufactured to give an optimal staining, when using the protocol advised in this IFU.

Prior to staining some routine fixed, paraffin-embedding tissue sections should be subjected to pre-treatment (HIER or digestive enzyme).

The BrightVision detection system detects Mouse/Rabbit bound to an antigen in tissue sections. The antibodies are not provided but it is recommended to use the WellMed-antibodies. This polymer-complex is then visualized with a suitable substrate/chromogen. The substrate is not provided but it is recommended to use the WellMed-substrate. The clinical interpretation of any staining or its absence should be determined by a qualified pathologist and complemented by morphologic studies; controls should be evaluated within the context of the patient's clinical history and/or other

3: Kit components

diagnostic tests.

BrightVision, one step detection system, Goat Anti- Mouse/Rabbit HRP (Ready-to-Use).







4: Availability:

Catalog Number	Contents	Volume
DPVO55HRP	BrightVision, one step detection system Goat Anti -Mouse/Rabbit HRP	55 ml
	(Ready-to-Use)	
DPVO110HRP	BrightVision, one step detection system Goat Anti -Mouse/Rabbit HRP	110 ml
	(Ready-to-Use)	
DPVO500HRP	BrightVision, one step detection system Goat Anti -Mouse/Rabbit HRP	500 ml
	(Ready-to-Use)	
DPVO999HRP	BrightVision, one step detection system Goat Anti -Mouse/Rabbit HRP	1000 ml
	(Ready-to-Use)	

5: Recommended Staining Protocol

Step	Reagent	Template step	Incubation time		
1	Deparaffinize and rehydrate tissue section	Slide/tissue preparing	-		
2	Wash Aqua dest	Wash	2x 5 min		
3	If applicable; HIER or digestive enzyme	Pre-treatment	*		
4	Wash buffer	PBS or TBS buffer	2x 5 min		
5	H ₂ O ₂ (concentration 3%)	Tissue preparing	10 min		
6	Wash buffer	PBS or TBS buffer	2x 5 min		
7	Primary rabbit or mouse antibody	Antibody	30 min		
8	Wash buffer	PBS or TBS buffer	2x 5 min		
9	Detection system, polymer Mouse/Rabbit HRP	Labeled polymer	30 min		
10	Wash buffer	PBS or TBS buffer	2x 5 min		
11	Substrate	DAB	*		
12	Wash aqua dest	Wash	2x 2 min		
13	Counterstain, dehydrate and coverslip	Auxiliary	-		
*					

* See applicable IFU

6: Control slides

A positive control, negative control and reagent control are needed and processed in the same way as the unknow specimen slide to interpret staining results.

7: Storage

Store at 2-8 °C and in the dark. Do not use after expiration date.

8: Warnings and precautions

Refer to SDS.

9: Troubleshooting

Please contact WellMed by phone or by email.

10: Reference

1) Shan-Rong Shi, James Guo, Richard J.cote, Lillian Young, Debra Hawes, Yan Shi, Sandra Thu and Clive R.Taylor, Applied Immunohistochemistry & Molecular Morphology, vol 7,201-208,1999

