

## MART-1 (Melan-A) clone M2-9E3


### Instructions for Use

**Specification:**

This antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This MAb labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive marker for angiomyolipoma's. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.

**Availability:**

Catalog No.	Contents	Volume
ILM2329-C01	MART-1 (Melan-A)	0,1 ml concentrate
ILM2329-C05	MART-1 (Melan-A)	0,5 ml concentrate
ILM2329-C1	MART-1 (Melan-A)	1,0 ml concentrate



**Intended use:** For Research Use Only

**Reactivity:** Human, Mouse and Rat. Others-not tested.

**Clone:** M2-9E3

**Species of origin:** Mouse

**Isotype:** IgG2b, kappa

**Control Tissue:** Melanoma, normal skin

**Staining:** Cytoplasmic

**Immunogen:** Recombinant hMART-1 protein

**Presentation:** Bioreactor Concentrate with 0.05% Azide

**Application and suggested dilutions:**

Pretreatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on formalin-fixed, paraffin embedded tissue sections.

- Immunohistochemical staining of cryostat tissue sections (dilution up to 1:50-1:100)
- Immunohistochemical staining of formalin-fixed, paraffin embedded tissue section (dilution up to 1:50-1:100)

The optimal dilution for a specific application should be determined by the investigator.

**Note:** Dilution of the antibody in 10% normal goat serum followed by a goat anti-mouse secondary antibody-based detection is recommended.

**Storage & Stability:** Store at 2-8 °C. Do not use after expiration date printed on the vial.

**References:**

- 1) Kawakami Y, *et. al.* Journal of Immunological Methods, 1997, 202(1):13-25.
- 2) Marincola FM, *et. al.* J of Immunotherapy with Emphasis on Tumor Immunol, 1996, 19(3):192-205.